



Effective mass of free neutrons in neutron star crust

Nicolas Chamel

*Copernicus Astronomical Center (CAMK), Polish Academy of Sciences, ul. Bartycka 18, 00-716 Warszawa, Poland
LUTH, Paris Observatory, 5 place Jules Janssen, 92195 Meudon, France*

Received 12 December 2005; received in revised form 20 March 2006; accepted 5 April 2006

Available online 8 June 2006

Abstract

The inner layers of a neutron star crust, composed of a Coulomb lattice of neutron rich nuclear clusters immersed in a sea of “free” superfluid neutrons, are closely analogous to periodic condensed matter systems such as electronic, photonic or phononic crystals. Applying methods from solid state physics to the neutron star context, it has been recently shown that Bragg scattering leads to a strong renormalization of the neutron mass in the bottom layers of the crust. Knowledge of this effective mass throughout the crust is essential in order to understand the dynamical properties of the neutron superfluid and the origin of pulsar glitches. The purpose of the present work is to evaluate this effective mass in the outermost layers of the inner crust, near the drip point $\rho_{\text{drip}} \sim 4 \times 10^{11} \text{ g cm}^{-3}$. The results are compared with the case of electrons in ordinary solids and as an example the corresponding effective electron mass in copper is calculated.

© 2006 Elsevier B.V. All rights reserved.

1. Introduction

The outer layers of a neutron star are formed of a solid crust composed of a Coulomb lattice of very neutron rich nuclei immersed in a nearly uniform relativistic electron gas. In the deeper layers corresponding to densities beyond the drip threshold $\rho_{\text{drip}} \simeq 4 \times 10^{11} \text{ g cm}^{-3}$, nuclei are embedded in a sea of “free” neutrons which are expected to become superfluids in mature neutron stars whose temperature has dropped below the critical temperature for the onset of superfluidity (for a review of neutron star crust matter, see [1] and references therein). Below the crust, at densities above $\sim 10^{14} \text{ g cm}^{-3}$ the “nuclei” merge into a uniform mixture of nucleons and leptons.

E-mail address: nchamel@camk.edu.pl (N. Chamel).

The crust of a neutron star, which represents only about 1% of the mass of the star and 10% of its radius, is nevertheless important in order to understand observational phenomena, such as for instance pulsar glitches. Pulsars are magnetized rotating neutron stars whose period, ranging from milliseconds to a few seconds, very slowly increases due to the loss of energy by electromagnetic and gravitational radiation. However some pulsars have been observed to suddenly spin up. These so-called “glitches” are characterized by a frequency jump, which varies from $\delta\Omega/\Omega \sim 10^{-9}$ (Crab) up to $\delta\Omega/\Omega \sim 10^{-6}$ (Vela), often accompanied by a permanent change in the slow down rate from $|\delta\dot{\Omega}/\dot{\Omega}| \sim 10^{-5}-10^{-4}$ (Crab) to $10^{-3}-10^{-2}$ (Vela). The relaxation following a glitch lasts from days to years. It was suggested by Baym et al. [2] soon after the discovery of the first pulsars that such very long time scales indicate the presence of superfluid components in the interior of the star. It is now widely accepted that pulsar glitches are due to a sudden transfer of angular momentum from the faster rotating neutron superfluid to the solid crust.

However the detailed mechanism of these glitches remain uncertain. In order to understand such phenomena, it is necessary to investigate the dynamical properties of the neutron superfluid in the inner crust. For this purpose, we have recently developed a simplified crust model in which the rigidity of the crust is neglected [3]. In this model, the crust is treated as a two fluid mixture: the neutron superfluid and the plasma of electrons and nuclei. The two fluids are assumed to be coupled by the so-called entrainment effects according to which the momentum of one fluid is a linear combination of the velocities of the two fluids [4]. This entrainment effect is completely determined by only one parameter \mathcal{K} from which an effective neutron mass m_* can be defined [3]. This effective mass is a crucial parameter for dynamical simulations of neutron stars and for understanding pulsar glitches [5]. For instance, it has been recently suggested that large enough effective masses could trigger a Kelvin–Helmholtz instability [6,7], which may provide a possible explanation for the origin of these glitches.

In a first series of work, we have calculated this effective mass in the very deep layers of the crust close to the interface with the liquid core in the “pasta” phases where the crust is formed by a lattice of slab (“lasagna”) or cylinder (“spaghetti”) shaped nuclei [8] and at lower densities for spherical nuclei [9]. We found that the effective mass is close to the bare neutron mass in the bottom of the crust while at lower densities the effective mass is strongly increased, reaching values as high as $m_* \sim 15m_n$ at the baryon density $5 \times 10^{13} \text{ g cm}^{-3}$. The purpose of the present work is to extend these calculations to much lower baryon densities of the order of $10^{11} \text{ g cm}^{-3}$ near the neutron drip point.

2. Band theory and effective mass

Since the classical paper of Negele & Vautherin [10] (subsequently referred simply as N&V), quantum calculations in the inner crust have been carried out in the Wigner–Seitz (W–S) approximation according to which the crust is decomposed into a compact arrangement of independent spheres centered around each nuclear cluster. This approach has been widely applied in the determination of the equilibrium structure of the crust and its static properties such as the equation of state for instance, since it greatly simplifies the numerical computations. Nevertheless the validity of this approach is questionable, in particular in the deep layers of the inner crust [11]. Furthermore since the nucleons are localized in the W–S sphere, this scheme cannot be applied for the study of transport properties and for the evaluation of the effective neutron mass. Since the “free” neutrons in the inner crust of a neutron star are closely analogous to “free” electrons in

metals, a natural step forward beyond the W–S approximation is to apply the well-known band theory.

This theory is not only restricted to the description of electrons in solids, but has been also applied to photonic and phononic crystals in which light (photons) and sound (phonons) respectively play the role of the electrons. In the band theory, the crust is assumed to be a perfect lattice whose composition is known. The clusters forming the crust are treated classically (this approximation is justified since each cluster contains about one hundred nucleons or more and are therefore much heavier than “free” neutrons) while the “free” neutrons are treated as independent particles subject to a periodic mean field [8]. We have recently shown how to extend this scheme in order to account for the pairing effects which are responsible for the superfluidity property [12]. By symmetry the problem can be reduced to the so-called W–S cell (a polyhedron) of the crystal, with the boundary conditions determined by the Floquet–Bloch theorem. In the W–S approximation, the cell is replaced by a sphere with more or less arbitrary boundary conditions (a detailed discussion can be found in Ref. [9]).

2.1. Effective neutron mass

In a previous work, we have shown that the effects of the pairing interaction on the effective mass (therefore on the entrainment parameters) are negligible when the spatial variations of the pairing field are small so that the BCS approximation is valid [12]. In the limit of an homogeneous system, the effective mass is independent of the pairing gap as confirmed by calculations of Gusakov & Haensel [13] in the liquid core of neutron stars. However it has been recently suggested that scattering on the inhomogeneities of the pairing field in the crust may be important and may even lead to localization of the “free” neutrons [14,15]. We shall here only consider the effective mass which results from normal Bragg scattering.

Subject to a more or less arbitrary convention specifying the density n_f of “free” neutrons as discussed in the previous works of this series [8,9], the effective neutron mass can then be expressed as

$$m_\star = \frac{n_f}{\mathcal{K}}. \tag{1}$$

The “mobility” coefficient \mathcal{K} is given by an integral over the Fermi surface

$$\mathcal{K} = \frac{1}{12\pi^3 \hbar^2} \sum_\alpha \int_{\text{F}} |\nabla_{\mathbf{k}} \mathcal{E}_{k\alpha}| dS^{(\alpha)}, \tag{2}$$

where $\mathcal{E}_{k\alpha}$ is the single particle energy of Bloch wave vector \mathbf{k} and band index α .

The effective mass m_\star , as defined by (1) and (2), has been referred in the solid state physics literature as an “optical” effective (electron) mass since it appears in the calculation of the real part of the dielectric constant of cubic metals [16], which can be written in the regime $\omega\tau \gg 1$ (τ being the relaxation time) as

$$\varepsilon\{\omega\} \simeq 1 - \frac{\omega_{p\star}^2}{\omega^2} + \varepsilon_{\text{inter}}, \tag{3}$$

where $\omega_{p\star}$ is an effective plasma frequency defined in terms of the density n_f of “free” electrons and of their effective mass m_\star by

$$\omega_{p\star}^2 = \frac{4\pi e^2 n_f}{m_\star} \tag{4}$$

and $\varepsilon_{\text{inter}}$ is the contribution due to interband transitions which is essentially constant for sufficiently low frequencies. It can be seen from (1) that this plasma frequency can be expressed as $\omega_{p\star}^2 = 4\pi e^2 \mathcal{K}$ and is therefore independent of the arbitrary convention used to specify which electrons are “free”.

It should also be remarked that the electric conductivity σ is related to the mobility coefficient \mathcal{K} introduced previously by the simple formula

$$\sigma = e^2 \tau \mathcal{K}, \quad (5)$$

where e is the electron electric charge and τ is the relaxation time. Unlike the effective electron mass m_\star , the electric conductivity (5) is obviously independent of the convention adopted for defining which electrons are considered to be “free”. This point has been usually somehow obscured in the literature where the conductivity is written in the form $\sigma = n_f e^2 \tau / m_\star$ of the classical Drude model.

This effective electron mass can therefore be deduced experimentally by studying the optical properties of solids or measuring the electric conductivity (or equivalently the resistivity $\rho = 1/\sigma$) provided the relaxation time τ is known. Values are typically in the range $m_\star \sim 1 - 2m_e$ for most elements except for beryllium which has a very small effective mass $m_\star/m_e = 0.46$ and lead whose estimates of effective mass reach $m_\star/m_e = 2.71$ as reported in Ref. [17] (see Appendix A for some numerical values of m_\star). The density n_f of “free” electrons is defined by counting for each atom in the crystal the number of electrons in the outermost atomic shell. For instance monovalent metals (such as alkali or noble metals) have only one “conduction” or “free” electron per atom in the crystal therefore $n_f = 1/\mathcal{V}_{\text{cell}}$ where $\mathcal{V}_{\text{cell}}$ is the volume of the W–S cell.

The mobility coefficient (2) can also be written as the integral over the Fermi volume

$$\mathcal{K} = \frac{1}{3} \sum_{\alpha} \int_{\text{F}} \text{Tr} \left\{ \frac{1}{m_\star\{\mathbf{k}\}^\alpha} \right\}, \quad (6)$$

of the trace of the *local* effective mass tensor [18] defined by

$$\frac{1}{m_\star\{\mathbf{k}\}^{\alpha_{ij}}} \equiv \frac{1}{\hbar^2} \frac{\partial^2 \mathcal{E}_{\alpha\mathbf{k}}}{\partial k_i \partial k_j}. \quad (7)$$

While the components of this tensor can be either positive or negative, it is clear from (2) that the “optical” effective mass defined by (1) is always positive.

The concept of the *local* effective mass tensor, which is well known in solid state physics, has been introduced only recently in the context of neutron diffraction [19]. The effective mass for neutrons propagating along Bragg planes is found to be very small compared to the bare mass by several orders of magnitude as expected from the analogy with electrons in solids (the electron effective mass at the edge of the band gap in semiconductors is reduced by about two orders of magnitude [18]). On the other hand, the “optical” effective mass m_\star as defined by (1) is an average of the local effective mass (7) over *all occupied states*, not only those for which the Bloch wave vector lies on a Bragg plane, and can therefore be expected to be much larger (it has already been pointed out that the electron effective mass is indeed usually increased in metals, see also Appendix A).

2.2. Band model of free neutrons

The structure of the inner crust has been determined by N&V within a density matrix expansion method whose equations are similar to those obtained with effective two body nucleon–

nucleon interactions of the Skyrme type in the Hartree–Fock approximation. We take their results for the lattice spacing and the composition of the nuclear clusters. The equilibrium structure of the crust is formed of a body centered cubic lattice.

From the self-consistent calculations of N&V, we construct an effective single particle equation for the “free” neutrons, neglecting spin–orbit coupling as was done by N&V (this approximation can be justified from the fact that the nuclear clusters, which are very neutron rich, have a very diffuse surface)

$$\hat{H}\varphi_{k\alpha} = \mathcal{E}_{k\alpha}\varphi_{k\alpha}, \tag{8}$$

$$\hat{H} = -\nabla \cdot \frac{\hbar^2}{2m_n^\oplus\{\mathbf{r}\}}\nabla + U_n\{\mathbf{r}\}. \tag{9}$$

As a result of the lattice symmetry, the single particle equations have to satisfy the Floquet–Bloch theorem [18]

$$\varphi_{k\alpha}\{\mathbf{r} + \mathbf{T}\} = e^{i\mathbf{k}\cdot\mathbf{T}}\varphi_{k\alpha}\{\mathbf{r}\}, \tag{10}$$

where \mathbf{T} is any lattice translation vector and \mathbf{k} is a wave vector belonging to the first Brillouin Zone (B-Z) of the reciprocal lattice. The equation can therefore be solved inside the W–S cell of the lattice (a truncated octahedron) with the boundary conditions (10). In the W–S approximation, this cell is replaced by a sphere with more or less arbitrary boundary conditions. For instance, N&V imposed the vanishing of the wave function or its radial derivative on the sphere depending on its parity. In this approximation, the \mathbf{k} dependence of the single particle states (resulting from the translational symmetry) is therefore neglected.

The kinetic term $-\hbar^2/2m_n^\oplus\{\mathbf{r}\}$ and the mean field potential $U_n\{\mathbf{r}\}$ are deduced from the self-consistent neutron and proton densities $n_n\{\mathbf{r}\}$ and $n_p\{\mathbf{r}\}$ obtained by N&V, by applying the extended Thomas–Fermi approximation up to second order with Skyrme interactions [20]

$$\begin{aligned} \frac{\hbar^2}{2m_n^\oplus\{\mathbf{r}\}} &= \frac{\hbar^2}{2m_n} + \frac{1}{8}(t_1(2+x_1) + t_2(2+x_2))n_b\{\mathbf{r}\} \\ &\quad - \frac{1}{8}(t_1(1+2x_1) - t_2(1+2x_2))n_n\{\mathbf{r}\}, \end{aligned} \tag{11}$$

$$\begin{aligned} U_n\{\mathbf{r}\} &= \frac{1}{2}t_0[(2+x_0)n_b\{\mathbf{r}\} - (1+2x_0)n_n\{\mathbf{r}\}] + \frac{1}{24}t_3[(2+x_3)(2+\gamma)n_b\{\mathbf{r}\}^{\gamma+1} \\ &\quad - (2x_3+1)(2n_b\{\mathbf{r}\}^\gamma n_n\{\mathbf{r}\} + \gamma n_b\{\mathbf{r}\}^{\gamma-1}(n_p\{\mathbf{r}\}^2 + n_n\{\mathbf{r}\}^2))] \\ &\quad + \frac{1}{8}[t_1(2+x_1) + t_2(2+x_2)]\tau\{\mathbf{r}\} + \frac{1}{8}[t_2(2x_2+1) - t_1(2x_1+1)]\tau_n\{\mathbf{r}\} \\ &\quad + \frac{1}{16}[t_2(2+x_2) - 3t_1(2+x_1)]\nabla^2 n_b\{\mathbf{r}\} \\ &\quad + \frac{1}{16}[3t_1(2x_1+1) + t_2(2x_2+1)]\nabla^2 n_n\{\mathbf{r}\}, \end{aligned} \tag{12}$$

where the kinetic densities are given by ($q = n, p$ for neutron and proton, respectively)

$$\tau_q = \tau_q^{\text{TF}} + \tau_q^{(2)}, \tag{13}$$

where τ_q^{TF} is the well-known Thomas–Fermi expression

$$\tau_q^{\text{TF}} = \frac{3}{5}(3\pi^2)^{2/3}n_q^{5/3} \tag{14}$$

and the second order contribution $\tau_q^{(2)} = \tau_q^{\text{BW}} + \tau_q^{\text{NL}}$ is given by the Bethe–Weizsäcker term

$$\tau_q^{\text{BW}} = \frac{1}{36} \frac{(\nabla n_q)^2}{n_q} + \frac{1}{3} \Delta n_q, \quad (15)$$

plus non-local terms arising from the Skyrme masses $f \equiv m/m_q^\oplus$ (calculated here for pure matter of the species q)

$$\tau_q^{\text{NL}} = \frac{1}{6} \frac{\nabla n_q \cdot \nabla f}{f} + \frac{1}{6} n_q \frac{\Delta f}{f} - \frac{1}{12} n_q \left(\frac{\nabla f}{f} \right)^2. \quad (16)$$

We have used the SkM [21] Skyrme force which is expected to give results close to those of N&V since its parameters like those in the density matrix expansion method of N&V, were adjusted on the neutron matter calculation of Siemens & Pandharipande [22] and since this force was used a few years later by Bonche & Vautherin [23] in the case of hot neutron star matter. For comparison, we have also used the more recent SLy4 force [24–26], which has been specifically constructed for astrophysical purposes and which has been already applied in neutron star crust calculations [27–32]. The parameters of these two Skyrme interactions are reported in Appendix C.

3. Numerical methods

3.1. LAPW method

We have solved Eq. (8) by the Linear Augmented Plane Wave (LAPW) method [33] which we have extended in order to allow for the presence of the space-dependent Skyrme mass $m_n^\oplus\{\mathbf{r}\}$. This method, whose idea was first introduced a long time ago by Slater [34] for the treatment of electrons in solids, provides a natural extension of the W–S approximation by combining the spherical symmetry around each nuclear cluster with the translational symmetry of the lattice.

The W–S cell is decomposed into two regions: a sphere of radius R_s surrounding the cluster and an interstitial region, where in the muffin tin approximation the Skyrme mass and the mean field are assumed to take uniform values.

Eq. (8) is solved by expanding the wave function on a mixed basis set

$$\varphi_{\mathbf{k}}\{\mathbf{r}\} = \sum_{\alpha} c_{\alpha} \phi_{\alpha\mathbf{k}}\{\mathbf{r}\}. \quad (17)$$

In the interstitial region the basis functions are defined by plane waves

$$\phi_{\alpha\mathbf{k}}\{\mathbf{r}\} = \frac{1}{\sqrt{V_{\text{cell}}}} e^{i(\mathbf{k} + \mathbf{K}_{\alpha}) \cdot \mathbf{r}}, \quad (18)$$

where \mathbf{K}_{α} are reciprocal lattice vectors, which ensures that the Floquet–Bloch theorem (10) is satisfied. Inside the Slater sphere, the basis functions are given by

$$\phi_{\alpha\mathbf{k}}\{\mathbf{r}\} = \sum_{\ell=0}^{\ell_{\text{max}}} \sum_{m=-\ell}^{\ell} (A_{\ell m} u_{\ell}\{\mathbf{r}\} + B_{\ell m} \dot{u}_{\ell}\{\mathbf{r}\}) Y_{\ell m}\{\hat{\mathbf{r}}\}, \quad (19)$$

where $Y_{\ell m}$ are spherical harmonics ($\hat{\mathbf{r}}$ refers to the angular coordinates of \mathbf{r}) and u_{ℓ} is the radial solution of Eq. (8) inside the sphere for a given energy \mathcal{E}_{ℓ} . The dot means derivative with respect to the energy. The coefficients $A_{\ell m}$ and $B_{\ell m}$ are determined by imposing the continuity of the wave function and of its gradient on the sphere (see [33] for further details).

From the Rayleigh–Ritz variational theorem, it follows that the Schrödinger equation (8) reduces to a generalized eigenvalue problem

$$\sum_{\beta} H_{\alpha\beta} c_{\beta} = \mathcal{E} \sum_{\beta} S_{\alpha\beta} c_{\beta}, \tag{20}$$

where $H_{\alpha\beta}$ is the matrix of the Hamiltonian (9)

$$H_{\alpha\beta} = \langle \phi_{\alpha k} | \hat{H} | \phi_{\beta k} \rangle, \tag{21}$$

and $S_{\alpha\beta}$ is the overlap matrix

$$S_{\alpha\beta} = \langle \phi_{\alpha k} | \phi_{\beta k} \rangle. \tag{22}$$

The expression of these matrices are given in Appendix B.

The expansion (17) is truncated such as to include all reciprocal lattice vectors satisfying

$$\frac{\hbar^2 (\mathbf{k} + \mathbf{K}_{\alpha})^2}{2m_n} < E_{\text{cutoff}}. \tag{23}$$

The choice of the radius R_s of the Slater sphere and of the linearization energies $\{\mathcal{E}_{\ell}\}$ is rather delicate due to the occurrence of “ghost” bands. These bands can be however easily identified from their low \mathbf{k} dependence (“flat” bands in the energy spectrum) and from their high sensitivity to small changes of the parameters R_s and \mathcal{E}_{ℓ} (for a discussion on this issue, see for instance [33]).

3.2. Fermi surface integration

Fermi surface integrations were carried out with the Gilat–Raubenheimer scheme [35]. The first B-Z is decomposed into small identical cubes, inside which the Fermi surface (if any) is approximated by a plane. From the knowledge of the energy bands and their gradient, it can be shown that the integral can be performed analytically inside each of those cubes. The integral over the whole Fermi surface is then obtained by summing the contributions of all the cubes. We have computed the energy gradient $\nabla_{\mathbf{k}} \mathcal{E}_{\mathbf{k}}$ by a finite difference scheme. It seems at first sight that the application of the Hellmann–Feynman theorem [36], from which the gradient is obtainable by the formula

$$\nabla_{\mathbf{k}} \mathcal{E}_{\mathbf{k}} = \sum_{\alpha, \beta} c_{\alpha}^* c_{\beta} (\nabla_{\mathbf{k}} H_{\alpha\beta} - \mathcal{E}_{\mathbf{k}} \nabla_{\mathbf{k}} S_{\alpha\beta}), \tag{24}$$

would lead to more accurate results. However this method, which we already applied in previous works [8,9], yield poor results in the present case because the errors in the wave functions obtained from the LAPW method are of order of $(\mathcal{E} - \mathcal{E}_{\ell})^2$, while the errors in the energy bands are only of order $(\mathcal{E} - \mathcal{E}_{\ell})^4$ [37].

We have tested the code by computing the energy bands and the effective mass m_{\star} in the empty lattice model [38]. For comparison, we have also computed the energy bands and “optical” effective electron mass of copper, which has been very often used as a benchmark for band structure calculation methods.

3.3. Application to copper

Copper is a noble metal whose electronic atomic configuration is $[\text{Ar}]3d^{10}4s^1$ ([Ar] denotes the electronic configuration of argon). A typical feature of this metal is the mixing or “hybridization” of s and d atomic shells which leads to significant distortions of the Fermi surface




Fig. 1. Fermi surface of copper obtained by Choy et al. [39] (available online at <http://www.phys.ufl.edu/fermisurface/>).

from a sphere as shown on Fig. 1 (in contrast to alkali metals for which the Fermi surface is nearly spherical).

The closed atomic shells $1s$, $2s$, $2p$, $3s$ and $3p$ give rise to fully occupied dispersionless energy bands in the crystal (“core” states) and it is therefore necessary to consider only the bands arising from the “valence” electrons in the $3d$ and $4s$ atomic shells. We have computed the energy bands of copper employing the classical Chodorov potential (see for instance [40] and references therein). Copper has a face centered cubic structure with a lattice spacing $a = 3.6147 \text{ \AA}$.

While the muffin-tin approximation is very good in neutron star crust, it is not so good in ordinary solids due to the long range of the Coulomb force. The radius R_s of the Slater sphere is thus set at the maximum possible value, namely half the nearest neighbor distance $R_s = \sqrt{2}a/4$. As already discussed by Koelling & Arbmán [37], the bands are rather sensitive to the choice of the linearization energies \mathcal{E}_ℓ due to the presence of narrow (“localized”) d bands with low \mathbf{k} dispersion. We thus calibrated the energies \mathcal{E}_ℓ by computing the band structure with the original non linear APW method [34]. This method is more accurate but is computationally much more expensive than the linearized version. The expansion of the basis functions was truncated to $E_{\text{cutoff}} = 15 \text{ Ryds}$ and $\ell_{\text{max}} = 4$. We found a fairly good agreement between the two methods with the following setting (energies are in Ryd): $\mathcal{E}_0 = -0.90$, $\mathcal{E}_1 = -0.20$, and $\mathcal{E}_2 = \mathcal{E}_3 = \mathcal{E}_4 = -0.70$. As can be seen on Fig. 2 the energy bands show some resemblance to the ideal Fermi spectrum except for low \mathbf{k} dispersion d bands in the range $-0.55 < \mathcal{E} < -0.8 \text{ Ryds}$. This is a characteristic feature of transition elements.

We have used the Chodorov potential to compute the mobility coefficient of copper and the associated electron (“optical”) effective mass according to Eqs. (1) and (2). For this calculation we have fixed all the linearization energies to the Fermi energy $\mathcal{E}_\ell = \mu = -0.385 \text{ Rydbergs}$. In solid state physics, the number of “free” electrons is usually taken as the number of electrons in the “conduction” band. In the case of copper, there is one “conduction” electron per ion in the crystal from the atomic $4s$ shell and therefore $n_f = 1/\mathcal{V}_{\text{cell}}$. The corresponding (“optical”) effective electron mass as defined by (1) and (2), $m_\star \simeq 1.285m_e$ obtained by integration over 11375 cubes in the irreducible domain (therefore 546000 cubes in the B-Z), is in agreement with the earlier estimate of Seagall [42] for the same potential $m_\star/m_e = 1.3 \pm 0.1$. There is however some ambiguities in the specification of the “free” electron density n_f in the case of transition metals since these elements may have different possible valences due to the presence of partially or completely filled d shell in the atom. We have also seen in the case of copper (see Fig. 2) that in the crystal the electrons from the $3d$ atomic shell strongly hybridize with the “conduction”




Fig. 2. Band structure of copper (energies are shown in Rydbergs) along high symmetry lines of the first B-Z (using the conventional labeling [41]), computed with the LAPW method (see the text for further details). For comparison we have also plotted the energy spectrum of the ideal Fermi gas suitably translated into the first B-Z. The horizontal dotted line indicates the position of the Fermi energy.

electrons from the outermost 4s shell. One may therefore adopt the convention of counting also these *d* electrons as “free”, in which case the effective mass becomes much larger $m_* \simeq 14.14m_e$ since $n_f = 11/\mathcal{V}_{\text{cell}}$.

4. Results and discussion

We have considered the largest W–S cell of N&V (the radius of the W–S sphere is $R_{\text{WS}} = 54$ fm) which corresponds to the onset of neutron drip at the baryon density $\rho = 4.66 \times 10^{11}$ g cm⁻³. The nucleon densities are expressed as Woods–Saxon functions inside the Slater sphere of radius R_s

$$n_q\{r\} = n_{q0} + (n_{qi} - n_{q0}) / (1 + e^{(r-R_q)/\xi_q}), \tag{25}$$

and are constants outside (and equal to $n_q\{R_s\}$). The parameters were adjusted so as to fit the density profiles given by N&V, imposing that $n_{po} = 0$ and $n_{no} = \tilde{\rho}_G$ where $\tilde{\rho}_G$ is the average density of the exterior neutron gas given by N&V. The lattice spacing a of the body centered cubic lattice was defined such that the volume of the W–S polyhedron, which is given by $\mathcal{V}_{\text{cell}} = a^3/2$ is equal to the volume of the W–S sphere, which leads to the formula

$$a = 2(\pi/3)^{1/3} R_{\text{WS}}. \tag{26}$$

We found good convergence with the energy cutoff $\mathcal{E}_{\text{cutoff}} = 1$ MeV and $\ell_{\text{max}} = 6$. We fixed the radius of the Slater sphere to $R_s = 35$ fm and all energy parameters \mathcal{E}_ℓ for $0 \leq \ell \leq \ell_{\text{max}}$ to the mid value 0.15 MeV for the range $0 < \mathcal{E} < 0.3$ MeV (the energy origin is chosen as the constant value of the potential outside the muffin tin). We compared the band structure with that of the non-linear APW method [34] and we found good agreement with this choice of parameters.

The single particle energy spectrum for the “free” neutrons is shown on Fig. 3 for wave vectors \mathbf{k} along specific symmetry lines of the first B-Z. The figure shows that, unlike the band structure of copper discussed in the previous section (see Fig. 2), the neutron energy spectrum is actually very close to that of an ideal Fermi gas, except for the presence of avoided crossings around some specific high symmetry points. The reason of this similarity lies in the so-called




Fig. 3. Energy bands (in MeV) of the “free” neutrons in neutron star crust (solid line) along high symmetry lines of the first B-Z (using the conventional labeling [41]) compared to the energy spectrum of the ideal Fermi gas (dashed line) suitably translated into the first B-Z for comparison. The band structure has been obtained with SkM Skyrme forces (details are explained in the text).

Phillips–Kleinman cancellation theorem [43]. As a result of the Pauli exclusion principle, the orthogonalization of the “free” neutron states to the “core” states leads to a repulsive non-local and energy-dependent potential which essentially vanishes outside nuclei and which partially compensates the strong attractive nuclear potential inside nuclei. In the present case, as the nuclei are very far apart ($R_n, R_p \ll R_{WS}$) and since the nuclear interactions are short range the screening is almost complete. Consequently the effects of the nuclear clusters on the “free” neutron gas are negligible for bulk properties involving all “free” or valence states. This is illustrated on Fig. 4 for the total energy density ε of the “free” neutrons given by

$$\varepsilon\{n_f\} = \int_0^\mu \mathcal{E} \mathcal{N}\{\mathcal{E}\} d\mathcal{E}, \quad n_f = \int_0^\mu \mathcal{N}\{\mathcal{E}\} d\mathcal{E}, \quad (27)$$

where $\mathcal{N}\{\mathcal{E}\}$ is the density of single particle states defined by an integral over the constant energy surface $\mathcal{E}_k = \mathcal{E}$

$$\mathcal{N}\{\mathcal{E}\} = \frac{2}{(2\pi)^3} \sum_\alpha \int \frac{dS_{\mathcal{E}}^{(\alpha)}}{|\nabla_{\mathbf{k}} \mathcal{E}_{k\alpha}|}. \quad (28)$$

It can be seen on Fig. 4 that the total energy density of the “free” neutrons does not depend much on the Skyrme interaction and is very close to that of the ideal Fermi gas. The Fermi surface integral (28) was computed for each energy on a grid of 20000 points in the range $0 < \mathcal{E} < 0.3$ MeV. Each of the Fermi surface integrations were performed with 9920 cubes in the irreducible domain, i.e. 476160 cubes in the first B-Z. Increasing the number of cubes did not lead to any noticeable change on the figure.

On the other hand, transport properties of the neutron gas depend only on the states lying on the Fermi surface whose shape is determined by the interactions between the clusters and the neutron gas (unlike its volume which is simply given by $\mathcal{V}_F = (2\pi)^3 n_n$ as in the ideal Fermi gas). As in the preceding work [9], the density n_f of “free” neutrons is defined by counting all single particle states whose energy is larger than the maximum value of the mean field potential $U_n\{\mathbf{r}\}$.




Fig. 4. Total energy density ε (in MeV per fm^3) of the “free” neutrons as a function of the “free” neutron density n_f (in fm^{-3}) compared with that of the ideal Fermi gas. The arrow indicates the “free” neutron density in the W–S cell considered by Negele & Vautherin.




Fig. 5. Effective neutron mass m_*/m_n as a function of the “free” neutron density n_f (in fm^{-3}). The arrow indicates the “free” neutron density in the W–S cell considered by Negele & Vautherin.

As can be seen on Fig. 5 the effective neutron mass m_* deviates significantly from the “bare” neutron mass m_n (as in the case of electrons in copper previously discussed in Section 3.3), and exhibits large variations with the density n_f due to band effects, except for very low dripped neutron densities n_f for which $m_* \sim m_n$.

This can be understood from the consideration that the effects of the nuclear clusters are very small whenever the Fermi wave length is much larger than the lattice spacing, or in other words when the Fermi volume lies well inside the first B-Z. This means that the Fermi surface is nearly spherical for densities much lower than the density for which the Fermi sphere touches the B-Z boundaries. Such a situation occurs for alkali metals. For a body centered cubic lattice, this density threshold is given by $n_f = \sqrt{2}\pi/3V_{\text{cell}}$ (at this density, the radius of the Fermi sphere equals the distance from the center of the first B-Z to the center of a face), which yields $n_f \simeq 2.2 \times 10^{-6} \text{ fm}^{-3}$ for $R_{\text{WS}} = 54 \text{ fm}$. Distortions of the Fermi sphere are expected




Fig. 6. Fermi surface area (in fm^{-2}) as a function of the “free” neutron density n_f (in fm^{-3}) compared to the area of the corresponding Fermi sphere. The arrow indicates the “free” neutron density in the W–S cell considered by Negele & Vautherin.

to occur for slightly lower densities due to the formation of necks as in copper (see Fig. 1). As shown on Fig. 6, above this density the Fermi surface area is reduced compared to the area of the corresponding sphere because by breaking the full translational symmetry (thereby raising some degeneracies), the presence of the nuclear lattice leads to the opening of holes on the Fermi surface. In the limit for which the translational symmetry is completely lost (as in the case of isolated nuclei or in the W–S approximation), the Fermi surface disappears and its area vanishes which obviously implies that $\mathcal{K} = 0$. The mobility coefficient can also vanish whenever the energy spectrum exhibits a gap at the Fermi level. This situation is realized in ordinary insulators and in semiconductors for which the electric conductivity $\sigma = e^2 \tau \mathcal{K} = 0$ since $\mathcal{K} = 0$ (at zero temperature). However this is still an open issue in the case of neutron star crust.

5. Conclusion

Extending previous works in the bottom layers of neutron star crust [8,9], we have applied the band theory of solids in order to evaluate the renormalization of the “free” neutron mass due to normal Bragg scattering in the outermost layers, at densities around $4 \times 10^{11} \text{ g cm}^{-3}$. The knowledge of this effective neutron mass is essential for the evaluation of the entrainment effects in hydrodynamical simulations of neutron stars [3,5]. In solid state physics, an analogous effective electron mass has been introduced in the study of the optical properties of materials. For comparison we have calculated the optical effective electron mass in copper and we have also reported values for other elements.

Using a phenomenological mean field model based on the results of Negele & Vautherin [10] with Bloch type boundary conditions, we found that the effects of the nuclear clusters are very small when the neutron gas is very dilute, meaning that the neutron Fermi wavelength is much larger than the lattice spacing. This is similar to what is observed in alkali metals. However at higher densities, the effective neutron mass m_* is significantly increased compared to the “bare” mass (as it is also observed for the case of electrons in ordinary metals) and fluctuates with the density as a result of band effects.

The present calculations performed in the outermost layers of the inner crust combined with those carried out in the bottom layers, suggest that the effective neutron mass m_*/m_n could take

very large values in the middle layers of the crust where the size of the nuclear clusters is of the same order as the lattice spacing. This raises the question whether in some layers the effective neutron mass could actually diverge due to the existence of band gaps in the energy spectrum. This question is one of the main issues in the study of any periodic materials such as for instance photonic or phononic crystals and have triggered considerable interests, both experimentally and theoretically. In our calculation, no such gaps were found for the outermost layers of the inner crust however as shown by Economou et al. [44] in the context of photonic crystals, this does not exclude the possibility of gaps at higher densities (higher energies), in deeper layers. The existence of band gaps would have important implications for the dynamics of the star and in particular for pulsar glitches.

Acknowledgements

The author acknowledges financial support from the Lavoisier program of the French Ministry of Foreign Affairs. The author is also very grateful to professor Haensel for discussions.

Appendix A. Effective electron mass in metals

Effective electron optical mass for a few elements taken from the compilation of Ref. [17].

Metal	m_*/m_e
Ag	0.99–1.04
Al	1.52
Au	1.1
Be	0.46
Ni	1.7
Ca	1.8
Cr	1.72
Fe	1.9
Cu	1.45
Ir	1.41
K	1.08
Li	1.45
Pb	2.12–2.71
Mo	1.43–2
Na	0.98–1.25
Nb	1.86
Pd	1.66
Rh	1.87
Ta	1.65
Ti	1.39
V	1.63
W	1.31

Appendix B. LAPW matrix elements

We have extended the LAPW method as discussed in the solid state context by Koelling and Arbman [37] in order to allow for a non-uniform mass in the kinetic term. The end result is simply the replacement of the electron mass m_e by the Skyrme neutron mass $m_n^\oplus\{R_s\}$ in the expression of the matrix elements. In Ref. [37], the authors used atomic units in which the kinetic factor $\hbar^2/2m_e$ is equal to unity and therefore the electron mass does not appear explicitly. For clarity and completeness, we report in this appendix the full expressions for the matrix elements.

The radial solution being normalized as

$$\int_0^{R_s} dr r^2 u_\ell\{r\}^2 = 1, \quad (\text{B.1})$$

the Hamiltonian and the overlap matrix elements are given by

$$\begin{aligned} \langle \phi_\alpha | \hat{H} \phi_\beta \rangle &= \frac{\hbar^2 \mathbf{q}_\alpha \cdot \mathbf{q}_\beta}{2m_n^\oplus\{R_s\}} \left(\delta_{\alpha\beta} - \frac{4\pi R_s^2}{\mathcal{V}_{\text{cell}}} \frac{j_1\{|\mathbf{q}_\alpha - \mathbf{q}_\beta\}}{|\mathbf{q}_\alpha - \mathbf{q}_\beta|} \right) \\ &+ \left(\frac{\hbar^2 R_s^2}{2m_n^\oplus\{R_s\}} \right)^2 \frac{4\pi}{\mathcal{V}_{\text{cell}}} \sum_{\ell=0}^{+\infty} (2\ell + 1) P_\ell\{\cos\theta\} [\mathcal{E}_\ell(a_\ell\{q_\alpha\}a_\ell\{q_\beta\} \\ &+ N_\ell b_\ell\{q_\alpha\}b_\ell\{q_\beta\}) + \gamma_{\alpha\beta}^\ell], \end{aligned} \quad (\text{B.2})$$

$$\begin{aligned} \langle \phi_\alpha | \phi_\beta \rangle &= \delta_{\alpha\beta} - \frac{4\pi R_s^2}{\mathcal{V}_{\text{cell}}} \frac{j_1\{|\mathbf{q}_\alpha - \mathbf{q}_\beta\}}{|\mathbf{q}_\alpha - \mathbf{q}_\beta|} \\ &+ \frac{4\pi R_s^2}{\mathcal{V}_{\text{cell}}} \sum_{\ell=0}^{+\infty} (a_\ell\{q_\alpha\}a_\ell\{q_\beta\} + N_\ell b_\ell\{q_\alpha\}b_\ell\{q_\beta\}), \end{aligned} \quad (\text{B.3})$$

where j_ℓ is the regular spherical Bessel function of order ℓ , P_ℓ is the Legendre polynomial of order ℓ , θ is the angle between the vectors $\mathbf{q}_\alpha \equiv \mathbf{k} + \mathbf{K}_\alpha$ and $\mathbf{q}_\beta \equiv \mathbf{k} + \mathbf{K}_\beta$ (where \mathbf{K}_α and \mathbf{K}_β are reciprocal lattice vectors),

$$N_\ell = \int_0^{R_s} dr r^2 \dot{u}_\ell\{r\}^2, \quad (\text{B.4})$$

$$a_\ell\{q\} = \dot{u}_\ell\{R_s\} j'_\ell\{q R_s\} - \dot{u}'_\ell\{R_s\} j_\ell\{q R_s\}, \quad (\text{B.5})$$

$$b_\ell\{q\} = u'_\ell\{R_s\} j_\ell\{q R_s\} - u_\ell\{R_s\} j'_\ell\{q R_s\}, \quad (\text{B.6})$$

$$\begin{aligned} \gamma_{\alpha\beta}^\ell &= \dot{u}_\ell\{R_s\} u'_\ell\{R_s\} (j'_\ell\{q_\alpha R_s\} j_\ell\{q_\beta R_s\} + j'_\ell\{q_\beta R_s\} j_\ell\{q_\alpha R_s\}) \\ &- \dot{u}'_\ell\{R_s\} u'_\ell\{R_s\} j_\ell\{q_\alpha R_s\} j_\ell\{q_\beta R_s\} - \dot{u}_\ell\{R_s\} u_\ell\{R_s\} j'_\ell\{q_\alpha R_s\} j'_\ell\{q_\beta R_s\}. \end{aligned} \quad (\text{B.7})$$

The prime means derivative with respect to the radial coordinate r .

Appendix C. Skyrme parameterizations

The values of the parameters of the Skyrme forces used in this work are given in the following table (the unit of energy is MeV and the unit of length is fm)

	SkM	SLy4
t_0	−2645	−2488.91
t_1	385	486.82
t_2	−120	−546.39
t_3	15595	13777
x_0	0.09	0.834
x_1	0	−0.3438
x_2	0	−1
x_3	0	1.354
γ	1/6	1/6

References

- [1] P. Haensel, Neutron star crusts, in: D. Blaschke, N.K. Glendenning, A. Sedrakian (Eds.), *Physics of Neutron Star Interiors*, Lecture Notes in Physics, vol. 578, Springer-Verlag, Berlin, 2001, pp. 127–174.
- [2] G. Baym, C. Pethick, D. Pines, M. Ruderman, Spin up in neutron stars: The future of the Vela pulsar, *Nature* 224 (1969) 872.
- [3] B. Carter, N. Chamel, P. Haensel, Entrainment coefficient and effective mass for conduction neutrons in neutron star crust: Macroscopic treatment, astro-ph/0408083, *Int. J. Mod. Phys.*, in press.
- [4] A.F. Andreev, E.P. Bashkin, Three velocity hydrodynamics of superfluid solutions, *Sov. Phys. JETP* 42 (1976) 164–167.
- [5] N. Chamel, B. Carter, Effect of entrainment on stress and pulsar glitches in stratified neutron star crust, *Mon. Not. R. Astron. Soc.* 368 (2006) 796–808.
- [6] N. Andersson, G.L. Comer, R. Prix, Are pulsar glitches triggered by a superfluid two-stream instability?, *Phys. Rev. Lett.* 90 (2003) 091101.
- [7] N. Andersson, G.L. Comer, R. Prix, The superfluid two-stream instability, *Mon. Not. R. Astron. Soc.* 354 (2004) 101–110.
- [8] B. Carter, N. Chamel, P. Haensel, Entrainment coefficient and effective mass for conduction neutrons in neutron star crust: Simple microscopic models, *Nucl. Phys. A* 748 (2005) 675–697.
- [9] N. Chamel, Band structure effects for dripped neutrons in neutron star crust, *Nucl. Phys. A* 747 (2005) 109–128.
- [10] J.W. Negele, D. Vautherin, Neutron star matter at sub-nuclear densities, *Nucl. Phys. A* 207 (1973) 298–320.
- [11] M. Baldo, E.E. Saperstein, S.V. Tolokonnikov, The role of the boundary conditions in the Wigner–Seitz approximation applied to the neutron star inner crust, nucl-th/0601096, *Phys. Lett. B*, in press.
- [12] B. Carter, N. Chamel, P. Haensel, Effect of BCS pairing on entrainment in neutron superfluid current in neutron star crust, *Nucl. Phys. A* 759 (2005) 164–441.
- [13] M. Gusakov, P. Haensel, The entrainment matrix of a superfluid neutron–proton mixture at a finite temperature, *Nucl. Phys. A* 761 (2005) 333–348.
- [14] A. Bulgac, P. Magierski, A. Wirzba, Fermionic Casimir effect in case of Andreev reflection, *Europhys. Lett.* 72 (2005) 327–333.
- [15] P. Magierski, Neutron localization induced by the pairing field in an inhomogeneous neutron matter, nucl-th/0508023.
- [16] M.H. Cohen, Optical constants, heat capacity, and the Fermi surface, *Philos. Mag.* 49 (1958) 762.
- [17] B. Hüttner, A new method for the determination of the optical mass of electrons in metals, *J. Phys.: Condens. Matter* 8 (1996) 11041–11052.
- [18] C. Kittel, *Introduction to Solid State Physics*, seventh ed., John Wiley & Sons, 1996.
- [19] A. Zeilinger, C.G. Shull, M.A. Horne, K. Finkelstein, Effective mass of neutrons diffracting in crystals, *Phys. Rev. Lett.* 57 (1986) 3089–3092.
- [20] M. Brack, C. Guet, H.B. Hakansson, Selfconsistent semiclassical description of average nuclear properties—a link between microscopic and macroscopic models, *Phys. Rep.* 123 (1985) 275–364.
- [21] H. Krivine, J. Treiner, O. Bohigas, Derivation of a fluid-dynamical Lagrangian and electric giant resonances, *Nucl. Phys. A* 336 (1980) 155–184.
- [22] P.J. Siemens, V.R. Pandharipande, Neutron matter computations in Bruckner and variational theories, *Nucl. Phys. A* 173 (1971) 561–570.

- [23] P. Bonche, D. Vautherin, A mean field calculation of the equation of state of supernova matter, Nucl. Phys. A 372 (1981) 496–526.
- [24] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, R. Schaeffer, New Skyrme effective forces for supernovae and neutron rich nuclei, Phys. Scr. T56 (1995) 231.
- [25] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, R. Schaeffer, A Skyrme parametrization from subnuclear to neutron star densities, Nucl. Phys. A 627 (1997) 710–746.
- [26] E. Chabanat, P. Bonche, P. Haensel, J. Meyer, R. Schaeffer, A Skyrme parametrization from subnuclear to neutron star densities Part II. Nuclei far from stabilities, Nucl. Phys. A 635 (1998) 231–256;
E. Chabanat, P. Bonche, P. Haensel, J. Meyer, R. Schaeffer, Nucl. Phys. A 643 (1998) 441, Erratum.
- [27] F. Douchin, P. Haensel, J. Meyer, Nuclear surface and curvature properties for SLy Skyrme forces and nuclei in the inner neutron star crust, Nucl. Phys. A 665 (2000) 419–446.
- [28] F. Douchin, P. Haensel, Inner edge of neutron-star crust with SLy effective nucleon–nucleon interactions, Phys. Lett. B 485 (2000) 107–114.
- [29] P. Magierski, A. Bulgac, P.-H. Heenen, Exotic nuclear phases in the inner crust of neutron stars in the light of the Skyrme–Hartree–Fock theory, Nucl. Phys. A 719 (2003) 217–220.
- [30] N. Sandulescu, N. van Giai, R.J. Liotta, Superfluid properties of the inner crust of neutron stars, Phys. Rev. C 69 (2004) 045802.
- [31] N. Sandulescu, Nuclear superfluidity and specific heat in the inner crust of neutron stars, Phys. Rev. C 70 (2004) 025801.
- [32] E. Khan, N. Sandulescu, N. van Giai, Collective excitations in the inner crust of neutron stars: Supergiant resonances, Phys. Rev. C 71 (2005) 042801.
- [33] D.J. Singh, Planewaves, Pseudopotentials and the LAPW Method, Kluwer Academic, Boston, 1994.
- [34] J.C. Slater, Wave functions in a periodic potential, Phys. Rev. 51 (1937) 846–851.
- [35] G. Gilat, L.J. Raubenheimer, Accurate numerical method for calculating frequency distribution functions in solids, Phys. Rev. 144 (2) (1966) 390–395;
G. Gilat, L.J. Raubenheimer, Accurate numerical method for calculating frequency distribution functions in solids, Phys. Rev. 147 (2) (1966) 670, Erratum.
- [36] R. Feynman, Forces in molecules, Phys. Rev. 56 (4) (1939) 340–343.
- [37] D.D. Koelling, G.O. Arbman, Use of energy derivative of the radial solution in an augmented plane wave method: Application to copper, J. Phys. F 5 (1975) 2041–2054.
- [38] W. Shockley, The empty lattice test of the cellular method in solids, Phys. Rev. 52 (1937) 866–872.
- [39] T.-S. Choy, J. Naset, J. Chen, S. Hershfield, C. Stanton, A database of Fermi surface in virtual reality modeling language (vrml), Bull. Am. Phys. Soc. 45 (2000) L36–L42.
- [40] G.A. Burdick, Energy band structure of copper, Phys. Rev. 129 (1963) 138–150.
- [41] G.F. Koster, in: F. Seitz, D. Turnbull (Eds.), Space Groups and Their Representations, Solid State Physics, vol. 5, Academic Press, New York, 1957.
- [42] B. Segall, Fermi surface and energy bands of copper, Phys. Rev. 125 (1962) 109–122.
- [43] J.C. Phillips, L. Kleinman, New method for calculating wave functions in crystals and molecules, Phys. Rev. 59 (1959) 287–294.
- [44] E.N. Economou, A. Zdetsis, Classical wave propagation in periodic structures, Phys. Rev. B 40 (1989) 1334–1337.